

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Terms	Documents
18 same nonionic	4

US Patents Full-Text Database

US Pre-Grant Publication Full-Text Database

JPO Abstracts Database

EPO Abstracts Database

Derwent World Patents Index

IBM Technical Disclosure Bulletins

Database:

18 same nonionic

Refine Search:

[Clear](#)**Search History**

Today's Date: 8/1/2001

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT	18 same nonionic	4	L10
USPT	15 same surfactant	0	L9
USPT	14 same surfactant	13	L8
USPT	14 same (hydrogenated near0 lecithin)	0	L7
USPT	15 same (hydrogenated near0 lecithin)	0	L6
USPT	14 same (growth near0 hormone\$)	10	L5
USPT	12 same l3	152	L4
USPT	mannitol same stabilit\$	888	L3
USPT	mannitol same stor\$	1009	L2
USPT	mannitol same (stor\$ and stabil\$)	512	L1

WEST**End of Result Set**

Generate Collection

L10: Entry 4 of 4

File: USPT

Dec 23, 1975

DOCUMENT-IDENTIFIER: US 3928137 A

TITLE: Reagent formulation for uric acid assay

ABPL:

An enzyme reagent formulation for use in assaying a biological specimen for uric acid is prepared as a granular, water-soluble, anhydrous, storage-stable mixture containing uricase, potassium chloride, mannitol, gum acacia, bovine serum albumin, glycine, sodium carbonate and a nitrogen containing polyoxyalkylene nonionic surfactant obtained by the sequential reaction of ethylenediamine with propylene oxide and ethylene oxide in the presence of a catalyst. The surfactant contains polyoxypropylene chains having an average molecular weight of between about 750 and about 6750, and polyoxyethylene chains constituting between about 10 and about 80 weight percent of the surfactant. The surfactant has an advantageous effect upon granulation, dissolution and storage stability of the reagent formulation.